



# Mathematical Inequalities & Applications

*With Compliments  
of the Author*

Zagreb, Croatia

Volume 25, Number 2, April 2022

*Gradimir V. Milovanović\* and Abdullah Mir*

*On the Erdős-Lax and Turán inequalities concerning  
polynomials*

# MATHEMATICAL INEQUALITIES & APPLICATIONS

## AIMS AND SCOPE

Mathematical Inequalities & Applications (MIA, Math. Inequal. Appl.) brings together original research papers in all areas of mathematics, provided they are concerned with inequalities or their role. From time to time MIA will publish invited survey articles. Short notes with interesting results or open problems will also be accepted.

MIA is published quarterly, in January, April, July and October.

## SUBMISSION

Manuscripts should be submitted electronically in PostScript or Adobe Acrobat PDF format to the MIA Editorial Office through MIA page: [www.ele-math.com](http://www.ele-math.com) or by e-mail [mia@ele-math.com](mailto:mia@ele-math.com)

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the MIA regarding the article. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

## COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to MIA. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal, and that its submission for publication has been approved by all the authors.

## PREPARATION OF MANUSCRIPT

Manuscripts should be written in English.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send final version of file(s) to the Editorial Office.

MIA is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

Mathematical Inequalities & Applications  
Editorial Office  
Menceticeva 2, 10000 Zagreb, Croatia  
[www.ele-math.com](http://www.ele-math.com)  
e-mail: [mia@ele-math.com](mailto:mia@ele-math.com)  
Fax: +385 1 6008799

## TITLE PAGE

A title page should include the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author and running head less than 60 characters. Authors should be listed alphabetically.

## ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2000) with primary (and secondary) subject classification codes and a list of 4-5 key words must be given.

## FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, grey scale only, preferably at resolution of 300 dpi or higher, sized to fit a journal page, and in EPS, TIFF or JPEG format. Figures should have a short descriptive title and should be numbered consecutively in order of appearance in the text. The position of each figure in the text should be indicated.

## REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The author should consult Mathematical Reviews for the standard abbreviations of journal names.

## PROOFS

The author will receive proofs only once. Late return of the proofs will delay the article to a later issue.

## PAGE CHARGES, OFFPRINTS

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from 'MIA' prior to publication.

## FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

## ON THE ERDŐS-LAX AND TURÁN INEQUALITIES CONCERNING POLYNOMIALS

GRADIMIR V. MILOVANOVIĆ\* AND ABDULLAH MIR

(Communicated by J. Jakšetić)

*Abstract.* In this paper, we prove certain sharp inequalities that relate the uniform norm of the derivative and the polynomial itself, in case when the zeros are outside or inside some closed disk. We further extend the obtained results to the polar derivative of a polynomial. The obtained results strengthen some recently proved Erdős-Lax and Turán-type inequalities contained in a paper published recently by Kumar [*Complex Anal. Oper. Theory* **14**, 65 (2020)], as well as other related inequalities.

### 1. Introduction

Let  $P(z)$  be a polynomial of degree  $n$  and  $P'(z)$  its derivative. Turán's inequality [23] that relates the norm of a polynomial to that of its derivative on the unit circle states that if  $P(z)$  is a polynomial of degree  $n$  having all its zeros in  $|z| \leq 1$ , then

$$\max_{|z|=1} |P'(z)| \geq \frac{n}{2} \max_{|z|=1} |P(z)|. \quad (1)$$

On the other hand, if  $P(z)$  has no zeros in  $|z| < 1$ , then Erdős conjectured and later Lax [10] proved that

$$\max_{|z|=1} |P'(z)| \leq \frac{n}{2} \max_{|z|=1} |P(z)|. \quad (2)$$

Thus in (1) as well as in (2) equality holds for those polynomials of degree  $n$  which have all their zeros on  $|z| = 1$ . Various versions of these inequalities are a classical topic in analysis. As a generalization of (1), Govil [7] proved that if  $P(z)$  has all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , then

$$\max_{|z|=1} |P'(z)| \geq \frac{n}{1+k^n} \max_{|z|=1} |P(z)|, \quad (3)$$

where as, for the class of polynomials not vanishing in  $|z| < k$ ,  $k \leq 1$ , the precise estimate of maximum of  $|P'(z)|$  on  $|z| = 1$  does not seem to be known in general. In

*Mathematics subject classification* (2020): 30A10, 30C10, 30D15.

*Keywords and phrases:* Polynomial, polar derivative, inequality, zeros.

\* Corresponding author.

1980, it was again Govil [6], who generalized (2) by proving that if  $P(z)$  does not vanish in  $|z| < k$ ,  $k \leq 1$ , then

$$\max_{|z|=1} |P'(z)| \leq \frac{n}{1+k^n} \max_{|z|=1} |P(z)|, \quad (4)$$

provided  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ , where  $Q(z) = z^n P(1/\bar{z})$ . Both the inequalities (3) and (4) are best possible and hold with equality for  $P(z) = z^n + k^n$ . Although the inequality (3) is sharp but it has a drawback. The bound in this inequality depends on the zero of the largest modulus and not on other zeros even if some of them are very close to the origin. This was taken into consideration by Aziz [1], who proved that if  $P(z) = a_n \prod_{v=1}^n (z - z_v)$  is a polynomial of degree  $n$  with  $|z_v| \leq k$ ,  $k \geq 1$ , then

$$\max_{|z|=1} |P'(z)| \geq \frac{2}{1+k^n} \sum_{v=1}^n \frac{k}{k+|z_v|} \max_{|z|=1} |P(z)|. \quad (5)$$

A similar type of modification to (4) was given by Aziz and Ahmad [2], who proved that if  $P(z) = a_n \prod_{v=1}^n (z - z_v)$  is a polynomial of degree  $n$  which does not vanish in  $|z| < k$ ,  $k \leq 1$  and  $Q(z) = z^n P(1/\bar{z})$ , then

$$\max_{|z|=1} |P'(z)| \leq \frac{1}{1+k^n} \left\{ n - k^n \sum_{v=1}^n \frac{|z_v| - k}{|z_v| + k} \right\} \max_{|z|=1} |P(z)|, \quad (6)$$

provided  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ . Equality in (6) holds for  $P(z) = z^n + k^n$ .

Very recently, Kumar [9] strengthened the bound in (5) by involving the modulus of each zero and some of the coefficients of the underlying polynomial. In fact, Kumar proved that if

$$P(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v)$$

is a polynomial of degree  $n$  having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , then

$$\max_{|z|=1} |P'(z)| \geq \left\{ \frac{2}{1+k^n} + \frac{(|a_n|k^n - |a_0|)(k-1)}{(1+k^n)(|a_n|k^n + k|a_0|)} \right\} \sum_{v=1}^n \frac{k}{|z_v| + k} \max_{|z|=1} |P(z)|. \quad (7)$$

In the same paper, Kumar proved an extension of (4) by showing that if  $P(z)$  is a polynomial of degree  $n$  having no zeros in  $|z| < k$ ,  $k \leq 1$ , and  $Q(z) = z^n P(1/\bar{z})$ , then

$$\max_{|z|=1} |P'(z)| \leq \left\{ n - \left[ \frac{2k^n}{1+k^n} + \frac{k^n(|a_0| - |a_n|k^n)(1-k)}{(1+k^n)(|a_0|k + |a_n|k^n)} \right] \sum_{v=1}^n \frac{|z_v|}{|z_v| + k} \right\} \max_{|z|=1} |P(z)|, \quad (8)$$

provided  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ . Equality in (7) and (8) holds for  $P(z) = z^n + k^n$ .

For a polynomial  $P(z)$  of degree  $n$ , we define

$$D_\alpha P(z) := nP(z) + (\alpha - z)P'(z),$$

the polar derivative of  $P(z)$  with respect to the point  $\alpha$ . The polynomial  $D_\alpha P(z)$  is of degree at most  $n - 1$  and it generalizes the ordinary derivative in the sense that

$$\lim_{\alpha \rightarrow \infty} \left\{ \frac{D_\alpha P(z)}{\alpha} \right\} = P'(z),$$

uniformly with respect to  $z$  for  $|z| \leq R$ ,  $R > 0$ .

Various results of majorization on the polar derivative of a polynomial can be found in the comprehensive books of Milovanović et al. [14], Marden [11] and Rahman and Schmeisser [22], where some approaches to obtaining polynomial inequalities are developed on applying the methods and results of the geometric function theory. For the latest research and development in this direction, one can see some of the papers ([9], [12], [13], [15]–[20]). By using the new version of the Schwarz lemma, Kumar [9] in the same paper also proved the polar derivative generalizations of (7) and (8) in the form of the following results.

**THEOREM A.** *Let  $P(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v)$  be a polynomial of degree  $n$  having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ . Then for any complex number  $\alpha$  with  $|\alpha| \geq k$ ,*

$$\begin{aligned} \max_{|z|=1} |D_\alpha P(z)| &\geq \frac{2(|\alpha| - k)}{1 + k^n} \left\{ 1 + \frac{(|a_n|k^n - |a_0|)(k - 1)}{2(|a_n|k^n + k|a_0|)} \right\} \\ &\quad \times \sum_{v=1}^n \frac{k}{|z_v| + k} \max_{|z|=1} |P(z)|. \end{aligned} \quad (9)$$

**THEOREM B.** *Let  $P(z) = \sum_{v=0}^n a_v z^v = a_n \prod_{v=1}^n (z - z_v)$  be a polynomial of degree  $n$  having no zeros in  $|z| < k$ ,  $k \leq 1$ , and  $Q(z) = z^n \overline{P(1/\bar{z})}$ . If  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ , then for any complex numbers  $\alpha$  with  $|\alpha| \geq 1$ ,*

$$\begin{aligned} \max_{|z|=1} |D_\alpha P(z)| &\leq \left\{ n|\alpha| - (|\alpha| - 1) \left[ \frac{2k^n}{1 + k^n} + \frac{k^n(|a_0| - |a_n|k^n)(1 - k)}{(1 + k^n)(|a_0|k + |a_n|k^n)} \right] \right\} \\ &\quad \times \sum_{v=1}^n \frac{|z_v|}{|z_v| + k} \max_{|z|=1} |P(z)|. \end{aligned} \quad (10)$$

**REMARK 1.** All the previous results can be formulated for monic polynomials, without loss of generality, so that we can put  $a_n = 1$  in the inequalities (7)–(10), obtaining simpler expressions.

If we divide both sides of (9) and (10) by  $|\alpha|$  and let  $|\alpha| \rightarrow \infty$ , we get respectively (7) and (8). It is important to mention here that Kumar [9] has mentioned in the last part of his paper that (8) and (10) are possibly the best available bounds so far towards the problem of generalizing the Erdős-Lax inequality for the class of polynomials having no zeros in  $|z| < k$ ,  $k \leq 1$ . Motivated by this, the authors are curious to establish some improved bounds of Erdős-Lax and Turán-type for the derivative and polar derivative of a polynomial. The obtained results produce refinements of (7)–(10) and related inequalities.

## 2. Main results

As we mentioned in Remark 1, without loss of generality, in the sequel we consider only the class of monic polynomials  $\widehat{\mathcal{P}}_n$  of degree  $n$ , with the complex zeros  $z_v$ ,  $v = 1, \dots, n$ , i.e.,

$$P(z) = z^n + a_{n-1}z^{n-1} + \dots + a_1z + a_0 = \prod_{v=1}^n (z - z_v), \quad (11)$$

for which we introduce the following quantities

$$A_n(k) = \sum_{v=1}^n \frac{k}{|z_v| + k}, \quad B_n(k) = \sum_{v=1}^n \frac{|z_v|}{|z_v| + k}, \quad m = \min_{|z|=k} |P(z)|. \quad (12)$$

We note that  $A_n(k) + B_n(k) = n$ .

We begin now by presenting the following strengthening of (7).

**THEOREM 1.** *Let  $P(z) \in \widehat{\mathcal{P}}_n$  having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ . Then for  $0 \leq t \leq 1$ , we have*

$$\begin{aligned} \max_{|z|=1} |P'(z)| &\geq \frac{2A_n(k)}{1+k^n} \left\{ \left[ 1 + \frac{(k^n - |a_0| - tm)(k-1)}{2(k^n + k|a_0| - tm)} \right] \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \frac{1}{2k^n} \left[ k^n - 1 - \frac{(k^n - |a_0| - tm)(k-1)}{k^n + k|a_0| - tm} \right] tm \right\}, \end{aligned} \quad (13)$$

Equality in (13) holds for  $P(z) = z^n + k^n$ .

**REMARK 2.** For  $t = 0$ , (13) reduces to (7) ( $a_n = 1$ ). In fact excepting the case when some or all the zeros of  $P(z)$  lie on  $|z| = k$ , the bound obtained in (13) is always sharper than the bound obtained in (7). As an illustration we consider two polynomials of degree four.

*Case (a):* Let  $P(z) = z^4 - 2z^3 + 4z - 4$ , with all zeros  $\{-\sqrt{2}, \sqrt{2}, 1 - i, 1 + i\}$  on the circle  $|z| = \sqrt{2}$ , so that Theorem 1 holds for  $k \geq \sqrt{2}$ . Since

$$M = \max_{|z|=1} |P(z)| = \max_{0 \leq \theta < 2\pi} \sqrt{37 - 36\cos\theta - 16\cos 2\theta + 24\cos 3\theta - 8\cos 4\theta}$$

and

$$M_1 = \max_{|z|=1} |P'(z)| = \max_{0 \leq \theta < 2\pi} \sqrt{68 - 48\cos\theta - 48\cos 2\theta + 32\cos 3\theta},$$

i.e.,  $M = 9.6142743738\dots$  and  $M_1 = 12.250756577\dots$ , as well as

$$m = \min_{|z|=k} |P(z)| = P(k) = (k^2 - 2)((k-1)^2 + 1) \quad (k \geq \sqrt{2}),$$

the right hand side of the inequality (13) is presented in Fig. 1 for  $t = 0$  and  $t = 1$ . The curve for  $t = 0$  is in fact, the right hand side of (7) ( $a_n = 1$ ).

*Case (b):* Let  $P(z) = z^4 + 4$ , with all zeros  $\{1 - i, 1 + i, -1 - i, -1 + i\}$  on the circle  $|z| = \sqrt{2}$ . Again Theorem 1 holds for  $k \geq \sqrt{2}$ , here with  $M = 5$ ,  $M_1 = 4$ , and  $m = \min_{|z|=k} |P(z)| = k^4 - 4$ . The corresponding graphics for  $t = 0$ ,  $t = 1/2$  and  $t = 1$  are presented in Fig. 2. As we can see the sharpest inequality is obtained for  $t = 1$ . Evidently, equality in (13) holds for  $k = \sqrt{2}$  for the polynomial  $z^4 + 4$ .




Figure 1: Bounds obtained by Theorem 1 when  $\sqrt{2} \leq k \leq 6$ : Case (a) for  $t = 0$  and  $t = 1$




Figure 2: Bounds obtained by Theorem 1 when  $\sqrt{2} \leq k \leq 6$ : Case (b) for  $t = 0, 1/2, 1$

It is also easy to see that (13) includes the following inequality due to Dewan and Upadhye [4]:

$$\max_{|z|=1} |P'(z)| \geq \sum_{v=1}^n \frac{k}{|z_v| + k} \left\{ \frac{2}{1+k^n} \max_{|z|=1} |P(z)| + \frac{k^n - 1}{k^n(k^n + 1)} \min_{|z|=k} |P(z)| \right\}.$$

In the sequel we prove the following refinement of (8), which in turn strengthens the bound in (6).

**THEOREM 2.** Let  $P(z) \in \widehat{\mathcal{P}}_n$  having no zeros in  $|z| < k$ ,  $k \leq 1$ , and let  $Q(z) = z^n \overline{P(1/\bar{z})}$ . If  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ , then for every  $0 \leq t \leq 1$ , we have

$$\begin{aligned} \max_{|z|=1} |P'(z)| &\leq \left\{ n - \left[ 1 + \frac{(|a_0| - k^n - tm)(1-k)}{2k(|a_0| + k^{n-1} - tm)} \right] \frac{2k^n B_n(k)}{1+k^n} \right\} \max_{|z|=1} |P(z)| \\ &\quad - tm \left( 1 - k^n - \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) \frac{B_n(k)}{1+k^n}. \end{aligned} \quad (14)$$

Equality in (14) holds for  $P(z) = z^n + k^n$ .

**REMARK 3.** For  $t = 0$ , (14) reduces to (8) ( $a_n = 1$ ). The following result which is a refinement of (6) immediately follows from Theorem 2 for  $t = 1$ .

**COROLLARY 1.** Let  $P(z) \in \widehat{\mathcal{P}}_n$  having no zeros in  $|z| < k$ ,  $k \leq 1$ , and let  $Q(z) = z^n \overline{P(1/\bar{z})}$ . If  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ , then

$$\begin{aligned} \max_{|z|=1} |P'(z)| &\leq \frac{1}{1+k^n} \left\{ \left( n - k^n \sum_{v=1}^n \frac{|z_v| - k}{|z_v| + k} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. - (1 - k^n) \sum_{v=1}^n \frac{|z_v|}{|z_v| + k} \min_{|z|=k} |P(z)| \right\}. \end{aligned} \quad (15)$$

Equality in (15) holds for  $P(z) = z^n + k^n$ .

In the literature, we can see a series of papers where inequalities involving the ordinary derivatives have been extended to the polar derivatives. Here, we are interested to extend Theorems 1 and 2 to the polar derivative of a polynomial and to get compact generalizations of these results. These generalizations involve the proofs of Theorems 1 and 2 as well, so it is obligatory to establish Theorems 1 and 2 before their respective generalizations in the form of Theorems 3 and 4. In this context, our next result is a polar derivative generalization of Theorem 1 which also provides a refinement of Theorem A.

**THEOREM 3.** Let  $P(z) \in \widehat{\mathcal{P}}_n$  having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ . Then for every complex number  $\alpha$  with  $|\alpha| \geq k$  and  $0 \leq t \leq 1$ , we have

$$\begin{aligned} \max_{|z|=1} |D_\alpha P(z)| &\geq \frac{2(|\alpha| - k)}{1+k^n} \left\{ \left( 1 + \frac{(k^n - |a_0| - tm)(k-1)}{2(k^n + k|a_0| - tm)} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \frac{1}{2k^n} \left( k^n - 1 - \frac{(k^n - |a_0| - tm)(k-1)}{k^n + k|a_0| - tm} \right) tm \right\} A_n(k). \end{aligned} \quad (16)$$

**REMARK 4.** For  $t = 0$ , Theorem 3 reduces to Theorem A ( $a_n = 1$ ). If we divide both sides of (16) by  $|\alpha|$  and let  $|\alpha| \rightarrow \infty$ , we get the inequality (13), and thus Theorem 3 is an extension of Theorem 1.

Finally, we prove the following polar derivative generalization of Theorem 2. The obtained inequality gives a refinement of the inequality (10) as well.

**THEOREM 4.** Let  $P(z) \in \widehat{\mathcal{P}}_n$  having no zeros in  $|z| < k$ ,  $k \leq 1$ , and let  $Q(z) = z^n \overline{P(1/\bar{z})}$ . If  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ , then for every complex number  $\alpha$  with  $|\alpha| \geq 1$  and  $0 \leq t \leq 1$ , we have

$$\begin{aligned} \max_{|z|=1} |D_\alpha P(z)| &\leq n |\alpha| \max_{|z|=1} |P(z)| \\ &- \frac{|\alpha|-1}{1+k^n} B_n(k) \left\{ 2k^n \left( 1 + \frac{(|a_0|-k^n-tm)(1-k)}{2k(|a_0|+k^{n-1}-tm)} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \left( 1 - k^n - \frac{k^{n-1}(|a_0|-k^n-tm)(1-k)}{|a_0|+k^{n-1}-tm} \right) tm \right\}. \end{aligned} \quad (17)$$

Equality in (17) holds for  $P(z) = z^n + k^n$ .

**REMARK 5.** For  $t = 0$ , Theorem 4 reduces to Theorem B ( $a_n = 1$ ). If we divide both sides of (17) by  $|\alpha|$  and let  $|\alpha| \rightarrow \infty$ , we get the inequality (14), and thus Theorem 4 is an extension of Theorem 2.

### 3. Auxiliary results

We need the following lemmas to prove our theorems. The following lemma is due to Mir et al. [20].

**LEMMA 1.** If  $P(z) \in \widehat{\mathcal{P}}_n$  having no zeros in  $|z| < 1$ , then for  $R \geq 1$  and  $0 \leq t \leq 1$ , we have

$$\begin{aligned} \max_{|z|=R} |P(z)| &\leq \left( \frac{(1+R^n)(|a_0|+R-tm_1)}{(1+R)(|a_0|+1-tm_1)} \right) \max_{|z|=1} |P(z)| \\ &- \left( \frac{(1+R^n)(|a_0|+R-tm_1)}{(1+R)(|a_0|+1-tm_1)} - 1 \right) tm_1, \end{aligned} \quad (18)$$

where  $m_1 = \min_{|z|=1} |P(z)|$ . Equality in (18) holds for  $P(z) = (\alpha + \beta z^n)/2$ ,  $|\alpha| = |\beta| = 1$ .

**LEMMA 2.** If  $P(z) \in \widehat{\mathcal{P}}_n$  having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , then for  $0 \leq t \leq 1$ , we have

$$\begin{aligned} \max_{|z|=k} |P(z)| &\geq \left( \frac{2k^n}{1+k^n} + \frac{k^n(k^n-|a_0|-tm)(k-1)}{(1+k^n)(k^n+k|a_0|-tm)} \right) \max_{|z|=1} |P(z)| \\ &+ \left( \frac{k^n-1}{k^n+1} - \frac{(k^n-|a_0|-tm)(k-1)}{(1+k^n)(k^n+k|a_0|-tm)} \right) tm, \end{aligned} \quad (19)$$

where  $m = \min_{|z|=k} |P(z)|$ . Equality in (19) holds for  $P(z) = z^n + k^n$ .

*Proof.* Let  $T(z) = P(kz)$ . Since  $P(z)$  has all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , the polynomial  $T(z)$  has all its zeros in  $|z| \leq 1$ . Let  $H(z) = z^n T(1/z)$  be the reciprocal

polynomial of  $T(z)$ , then  $H(z)$  is a polynomial of degree at most  $n$  having no zeros in  $|z| < 1$ . Hence applying (18) of Lemma 1 to the polynomial  $H(z)$ , we get for  $k \geq 1$  and  $0 \leq t \leq 1$ ,

$$\begin{aligned} \max_{|z|=k} |H(z)| &\leq \left( \frac{(1+k^n)(k^n + k|a_0| - tm^*)}{(1+k)(k^n + |a_0| - tm^*)} \right) \max_{|z|=1} |H(z)| \\ &\quad - \left( \frac{(1+k^n)(k^n + k|a_0| - tm^*)}{(1+k)(k^n + |a_0| - tm^*)} - 1 \right) tm^*, \end{aligned} \quad (20)$$

where  $m^* = \min_{|z|=1} |H(z)|$ .

Since  $|H(z)| = |T(z)|$  on  $|z| = 1$ , hence,

$$\begin{aligned} m^* &= \min_{|z|=1} |H(z)| = \min_{|z|=1} \left| z^n P\left(\frac{k}{z}\right) \right| = \min_{|z|=k} |P(z)| = m, \\ \max_{|z|=1} |H(z)| &= \max_{|z|=1} |T(z)| = \max_{|z|=k} |P(z)|, \end{aligned}$$

and

$$\max_{|z|=k} |H(z)| = \max_{|z|=k} \left| z^n P\left(\frac{k}{z}\right) \right| = k^n \max_{|z|=1} |P(z)|,$$

which when substituted in (20) gives

$$\begin{aligned} \max_{|z|=k} |P(z)| &\geq \left( \frac{(1+k)(k^n + |a_0| - tm)}{(1+k^n)(k^n + k|a_0| - tm)} \right) k^n \max_{|z|=1} |P(z)| \\ &\quad + \left( 1 - \frac{(1+k)(k^n + |a_0| - tm)}{(1+k^n)(k^n + k|a_0| - tm)} \right) tm. \end{aligned} \quad (21)$$

Using the fact that

$$\frac{(1+k)(k^n + |a_0| - tm)}{(1+k^n)(k^n + k|a_0| - tm)} = \frac{2}{1+k^n} + \frac{(k^n - |a_0| - tm)(k-1)}{(1+k^n)(k^n + k|a_0| - tm)},$$

in (21), we get

$$\begin{aligned} \max_{|z|=k} |P(z)| &\geq \left( \frac{2k^n}{1+k^n} + \frac{k^n(k^n - |a_0| - tm)(k-1)}{(1+k^n)(k^n + k|a_0| - tm)} \right) \max_{|z|=1} |P(z)| \\ &\quad + \left( \frac{k^n - 1}{k^n + 1} - \frac{(k^n - |a_0| - tm)(k-1)}{(1+k^n)(k^n + k|a_0| - tm)} \right) tm, \end{aligned}$$

which is (19) and this completes the proof of Lemma 2.  $\square$

LEMMA 3. If  $P(z) \in \widehat{\mathcal{P}}_n$  and  $Q(z) = z^n \overline{P(1/\bar{z})}$ , then on  $|z| = 1$ ,

$$|P'(z)| + |Q'(z)| \leq n \max_{|z|=1} |P(z)|.$$

The above lemma is due to Govil and Rahman [8].

LEMMA 4. If  $P(z)$  is a polynomial of degree  $n$ , then for  $R \geq 1$ ,

$$\max_{|z|=R} |P(z)| \leq R^n \max_{|z|=1} |P(z)|.$$

The above lemma is a simple consequence of the Maximum Modulus Principle [21]. The following lemma is due to Giroux, Rahman and Schmeisser [5].

**LEMMA 5.** *If  $P(z) = \prod_{v=1}^n (z - z_v)$  is a polynomial of degree  $n$  having all its zeros in  $|z| \leq 1$ , then*

$$\max_{|z|=1} |P'(z)| \geq \sum_{v=1}^n \frac{1}{1 + |z_v|} \max_{|z|=1} |P(z)|.$$

#### 4. Proofs of main results

*Proof of Theorem 1.* Let  $G(z) = P(kz)$ . Recall that a monic polynomial  $P(z)$  has all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , the polynomial

$$G(z) = k^n \prod_{v=1}^n \left( z - \frac{z_v}{k} \right)$$

has all its zeros in  $|z| \leq 1$ , therefore, applying Lemma 5 to  $G(z)$ , we get

$$\max_{|z|=1} |G'(z)| \geq \sum_{v=1}^n \frac{1}{1 + \frac{|z_v|}{k}} \max_{|z|=1} |G(z)|, \quad (22)$$

which is equivalent to

$$k \max_{|z|=1} |P'(kz)| \geq \sum_{v=1}^n \frac{k}{k + |z_v|} \max_{|z|=k} |P(z)|. \quad (23)$$

Since  $P'(z)$  is a polynomial of degree  $n-1$  and  $k \geq 1$ , therefore by Lemma 4, we have

$$\max_{|z|=1} |P'(kz)| = \max_{|z|=k} |P'(z)| \leq k^{n-1} \max_{|z|=1} |P'(z)|.$$

Using this and Lemma 2 in (23), as well as the notation from (12), we get

$$\begin{aligned} k^n \max_{|z|=1} |P'(z)| &\geq \frac{A_n(k)}{1+k^n} \left\{ \left( 2k^n + \frac{k^n(k^n - |a_0| - tm)(k-1)}{k^n + k|a_0| - tm} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \left( k^n - 1 - \frac{(k^n - |a_0| - tm)(k-1)}{k^n + k|a_0| - tm} \right) tm \right\}, \end{aligned}$$

which is equivalent to (13). This completes the proof of Theorem 1.  $\square$

*Proof of Theorem 2.* Let  $Q(z) = z^n \overline{P(1/\bar{z})}$ . Since  $P(z) \neq 0$  in  $|z| < k$ ,  $k \leq 1$ , the polynomial  $Q(z)$  of degree  $n$  has all its zeros in  $|z| \leq 1/k$ ,  $1/k \geq 1$ . On applying

Theorem 1 to  $Q(z)$ , we get for  $0 \leq t \leq 1$ ,

$$\begin{aligned} \max_{|z|=1} |Q'(z)| &\geq \sum_{v=1}^n \frac{\frac{1}{k}}{\frac{1}{k} + \frac{1}{|z_v|}} \left[ \left( \frac{2}{1 + \frac{1}{k^n}} + \frac{\left(\frac{|a_0|}{k^n} - 1 - \frac{tm}{k^n}\right) \left(\frac{1}{k} - 1\right)}{\left(1 + \frac{1}{k^n}\right) \left(\frac{|a_0|}{k^n} + \frac{1}{k} - \frac{tm}{k^n}\right)} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \left( \frac{\frac{1}{k^n} - 1}{\frac{1}{k^n} \left(\frac{1}{k^n} + 1\right)} - \frac{\left(\frac{|a_0|}{k^n} - 1 - \frac{tm}{k^n}\right) \left(\frac{1}{k} - 1\right)}{\frac{1}{k^n} \left(\frac{1}{k^n} + 1\right) \left(\frac{|a_0|}{k^n} + \frac{1}{k} - \frac{tm}{k^n}\right)} \right) \frac{tm}{k^n} \right], \end{aligned} \quad (24)$$

because

$$\min_{|z|=1/k} |Q(z)| = \min_{|z|=1/k} \left| z^n \overline{P(1/\bar{z})} \right| = \frac{1}{k^n} \min_{|z|=k} |P(z)| = \frac{m}{k^n}$$

and

$$\max_{|z|=1} |Q(z)| = \max_{|z|=1} |P(z)|.$$

The above inequality (24) is equivalent to

$$\begin{aligned} \max_{|z|=1} |Q'(z)| &\geq \frac{B_n(k)}{1+k^n} \left\{ \left( 2k^n + \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \left( 1 - k^n - \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) tm \right\}, \end{aligned} \quad (25)$$

where  $B_n(k)$  is defined in (12).

Also  $|P'(z)|$  and  $|Q'(z)|$  attain maximum at the same point on  $|z| = 1$ , we have

$$\max_{|z|=1} (|P'(z)| + |Q'(z)|) = \max_{|z|=1} |P'(z)| + \max_{|z|=1} |Q'(z)|. \quad (26)$$

On combining (25), (26) and Lemma 3, we get

$$\begin{aligned} n \max_{|z|=1} |P(z)| &\geq \max_{|z|=1} |P'(z)| \\ &\quad + \frac{B_n(k)}{1+k^n} \left\{ \left( 2k^n + \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \left( 1 - k^n - \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) tm \right\}, \end{aligned}$$

which after simplification yields (14).

This proves Theorem 2 completely.  $\square$

*Proof of Theorem 3.* Let  $G(z) = P(kz)$ . Since  $P(z)$  has all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , therefore, all the zeros of  $G(z)$  lie in  $|z| \leq 1$ , hence by (22), we get

$$\max_{|z|=1} |G'(z)| \geq \sum_{v=1}^n \frac{k}{k + |z_v|} \max_{|z|=1} |G(z)|. \quad (27)$$

Let  $H(z) = z^n \overline{G(1/\bar{z})}$ . Then it can be easily verified that

$$|H'(z)| = |nG(z) - zG'(z)| \text{ for } |z| = 1. \quad (28)$$

The polynomial  $H(z)$  has all its zeros in  $|z| \geq 1$  and  $|H(z)| = |G(z)|$  for  $|z| = 1$ , therefore, by the result of de-Bruijn [3],

$$|H'(z)| \leq |G'(z)| \text{ for } |z| = 1. \quad (29)$$

Now, noting that by hypothesis, we have  $|\alpha|/k \geq 1$ , hence on using definition of polar derivative of a polynomial, we get

$$\begin{aligned} |D_{\alpha/k}G(z)| &= \left| nG(z) + \frac{\alpha}{k}G'(z) - zG'(z) \right| \\ &\geq \left| \frac{\alpha}{k} \right| |G'(z)| - |nG(z) - zG'(z)|, \end{aligned}$$

which on using (28) and (29), gives

$$\left| D_{\frac{\alpha}{k}}G(z) \right| \geq \frac{|\alpha|-k}{k} \max_{|z|=1} |G'(z)|. \quad (30)$$

Using (27) in (30) and on replacing  $G(z)$  by  $P(kz)$ , we get

$$\max_{|z|=1} |D_{\alpha/k}P(kz)| \geq \frac{|\alpha|-k}{k} \sum_{v=1}^n \frac{k}{k+|z_v|} \max_{|z|=1} |P(kz)|,$$

which implies

$$\max_{|z|=1} \left| nP(kz) + \left( \frac{\alpha}{k} - z \right) kP'(kz) \right| \geq \frac{|\alpha|-k}{k} \sum_{v=1}^n \frac{k}{k+|z_v|} \max_{|z|=k} |P(z)|,$$

which gives by using Lemma 2 on the right hand side that

$$\begin{aligned} \max_{|z|=k} |D_{\alpha}P(z)| &\geq \frac{|\alpha|-k}{k} \frac{A_n(k)}{1+k^n} \left\{ \left( 2k^n + \frac{k^n(k^n - |a_0| - tm)(k-1)}{k^n + k|a_0| - tm} \right) \max_{|z|=1} |P(z)| \right. \\ &\quad \left. + \left( k^n - 1 - \frac{(k^n - |a_0| - tm)(k-1)}{k^n + k|a_0| - tm} \right) tm \right\}, \end{aligned} \quad (31)$$

where  $A_n(k)$  is defined in (12). Since  $D_{\alpha}P(z)$  is a polynomial of degree at most  $n-1$  and  $k \geq 1$ , applying Lemma 4 to the polynomial  $D_{\alpha}P(z)$ , we get

$$\max_{|z|=k} |D_{\alpha}P(z)| \leq k^{n-1} \max_{|z|=1} |D_{\alpha}P(z)|,$$

which on using in (31) gives (16).

This completes the proof of Theorem 3.  $\square$

*Proof of Theorem 4.* Since  $Q(z) = z^n \overline{P(1/\bar{z})}$ , it is easy to verify that for  $|z| = 1$ ,

$$|Q'(z)| = |nP(z) - zP'(z)|. \quad (32)$$

Also, for any complex number  $\alpha$  with  $|\alpha| \geq 1$ , the polar derivative of  $P(z)$  with respect to  $\alpha$  is

$$D_{\alpha}P(z) = nP(z) + (\alpha - z)P'(z).$$

This implies by (32) for  $|z| = 1$ , that

$$\begin{aligned} |D_\alpha P(z)| &\leq |nP(z) - zP'(z)| + |\alpha||P'(z)| \\ &= |Q'(z)| + |\alpha||P'(z)| \\ &= |Q'(z)| + |P'(z)| - |P'(z)| + |\alpha||P'(z)| \\ &\leq n \max_{|z|=1} |P(z)| + (|\alpha| - 1)|P'(z)| \quad (\text{by Lemma 3}). \end{aligned}$$

This gives by using Theorem 2, that

$$\begin{aligned} \max_{|z|=1} |D_\alpha P(z)| &\leq n \max_{|z|=1} |P(z)| \\ &+ (|\alpha| - 1) \left[ \left\{ n - \left( 2k^n + \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) \frac{B_n(k)}{1+k^n} \right\} \max_{|z|=1} |P(z)| \right. \\ &\quad \left. - tm \left( 1 - k^n - \frac{k^n(|a_0| - k^n - tm)(1-k)}{k(|a_0| + k^{n-1} - tm)} \right) \frac{B_n(k)}{1+k^n} \right], \end{aligned}$$

which after simplification gives (17).

This completes the proof of Theorem 4.

## 5. Conclusion

We proved some sharp inequalities that relate the uniform norm of the derivative and the polynomial itself, in case when the zeros are outside or inside some closed disk. Also, we extended these results to the polar derivative of a polynomial. The obtained results sharpen and generalize some already known estimates of Erdős-Lax and Turán-type.

*Acknowledgements.* The first author was partly supported by the Serbian Academy of Sciences and Arts (Project Φ-96). The authors are grateful to the referees for carefully reading the manuscript, finding some mistakes and making certain useful suggestions.

## REFERENCES

- [1] A. AZIZ, *Inequalities for the derivative of a polynomial*, Proc. Amer. Math. Soc., **89** (1983), 259–266.
- [2] A. AZIZ AND N. AHMAD, *Inequalities for the derivative of a polynomial*, Proc. Indian Acad. Sci. (Math. Sci.), **107** (1997), 189–196.
- [3] N. G. DE BRUIJN, *Inequalities concerning polynomials in the complex domain*, Nederl. Akad. Wetensch Proc., **50** (1947), 1265–1272.
- [4] K. K. DEWAN AND C. M. UPADHYE, *Inequalities for the polar derivative of a polynomial*, J. Ineq. Pure Appl. Math., **9** (2008), Art. 119, pp. 1–9.
- [5] A. GIROUX, Q. I. RAHMAN AND G. SCHMEISSER, *On Bernstein's inequality*, Canad. J. Math., **31** (1979), 347–353.
- [6] N. K. GOVIL, *On a theorem of S. Bernstein*, Proc. Nat. Acad. Sci., **50** (1980), 50–52.
- [7] N. K. GOVIL, *On the derivative of a polynomial*, Proc. Amer. Math. Soc., **41** (1973), 543–546.

- [8] N. K. GOVIL AND Q. I. RAHMAN, *Functions of exponential type not vanishing in a half plane and related polynomials*, Trans. Amer. Math. Soc., **137** (1969), 501–517.
- [9] P. KUMAR, *On the inequalities concerning polynomials*, Complex Anal. Oper. Theory, **14**: 65 (2020), <https://doi.org/10.1007/s11785-020-01023-0>.
- [10] P. D. LAX, *Proof of a conjecture of P. Erdős on the derivative of a polynomial*, Bull. Amer. Math. Soc., **50** (1944), 509–513.
- [11] M. MARDEN, *Geometry of Polynomials*, Math. Surveys, No. 3, Amer. Math. Soc., Providence, R.I., 1966.
- [12] G. V. MILOVANOVIĆ AND A. MIR, *On the Erdős-Lax inequality concerning polynomials*, Math. Inequal. Appl. **23** (2020), 1499–1508.
- [13] G. V. MILOVANOVIĆ AND A. MIR, *Generalizations of Zygmund-type integral inequalities for the polar derivative of a complex polynomial*, J. Inequal. Appl. 2020, paper no. 136, 12 pp.
- [14] G. V. MILOVANOVIĆ, D. S. MITRINOVIĆ AND TH. M. RASSIAS, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific, Singapore, 1994.
- [15] A. MIR, *Bernstein-type integral inequalities for a certain class of polynomials*, Mediterranean J. Math., **16** (2019), Art. 143, pp. 1–11.
- [16] A. MIR, *Generalizations of Bernstein and Turán-type inequalities for the polar derivative of a complex polynomial*, Mediterranean J. Math., **17** (2020), Art. 14, pp. 1–12.
- [17] A. MIR, *On an operator preserving inequalities between polynomials*, Ukrainian Math. J., **69** (2018), 1234–1247.
- [18] A. MIR AND I. HUSSAIN, *On the Erdős-Lax inequality concerning polynomials*, C. R. Acad. Sci. Paris Ser. I, **355** (2017), 1055–1062.
- [19] A. MIR AND A. WANI, *A note on two recent results about polynomials with restricted zeros*, J. Math. Inequal., **14** (2020), 47–52.
- [20] A. MIR, I. HUSSAIN AND A. WANI, *A note on Ankeny-Rivlin theorem*, J. Anal., **27** (2019), 1103–1107.
- [21] G. PÓLYA AND G. SZEGŐ, *Aufgaben und Lehrsätze aus der Analysis*, Springer, Berlin, 1981.
- [22] Q. I. RAHMAN AND G. SCHMEISSER, *Analytic Theory of Polynomials*, Oxford University Press Inc. New York, 2002.
- [23] P. TURÁN, *Über die Ableitung von polynomen*, Compositio Math., **7** (1939), 89–95.

(Received May 24, 2021)

Gradimir V. Milovanović  
Serbian Academy of Sciences and Arts  
Knez Mihailova 35, 11000 Belgrade, Serbia  
and  
University of Niš  
Faculty of Science and Mathematics  
Niš, Serbia  
e-mail: gvm@mi.sanu.ac.rs

Abdullah Mir  
Department of Mathematics  
University of Kashmir  
190006 Srinagar, India  
e-mail: drabmir@yahoo.com

